arXiv:quant-ph/0411095v1 13 Nov 2004

Non-decomposable Quantum Dynamical Semigroups and Bound Entangled States

Fabio Benatti
Dipartimento di Fisica Teorica, Università di Trieste & Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, Trieste, Italy

Roberto Floreanini
Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, Trieste, Italy

Marco Piani
Dipartimento di Fisica Teorica, Università di Trieste & Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, Trieste, Italy

Abstract. We use open quantum system techniques to construct one-parameter semigroups of positive maps and apply them to study the entanglement properties of a class of 16-dimensional density matrices, representing states of a 4×4 bipartite system.

1. Introduction

The recent developments in quantum information have turned entanglement into a concrete physical resource that is important to identify, quantify and classify (*e.g.* see [1, 2] and references therein). The techniques devised to this end have spurred the novel use of some mathematical notions like positive linear maps [3–5] which had been somewhat neglected in quantum theory; in fact, fully consistent physical operations have to be described by completely positive maps [6–8].

The need for complete positivity is due to the existence of entangled states: entangled bipartite states, if subjected to positive transformations acting on one partner only, do not in general remain positive and thus lose their probabilistic interpretation (for recent related work on this issue, see [9–13]). However, this very same fact renders positive maps useful in detecting entanglement [14–18].

The simplest instance of positive, but not completely positive map, is the transposition: when acting on one partner only (partial transposition) of a bipartite system it detects all entangled states of 2×2 and 2×3-dimensional systems. In higher dimension, however, there are bipartite states which remain positive under partial transposition and are nevertheless entangled. Further, the kind of entanglement they contain cannot be distilled by local quantum operations and classical communication (these and other properties are reviewed *e.g.* in [18]).

This undistillable entanglement is termed “bound” and can be detected only by means of indecomposable positive maps, namely maps that cannot be written as the sum of a completely positive map and a completely positive map composed with transposition. It is thus of practical importance to provide as many examples of indecomposable maps as possible, but the task is rather difficult for, unlike completely positive maps, the structure of positive ones is still elusive [3, 4].

In this work, we add to the phenomenology of positive maps [19–21] a class of one-parameter semigroups [22] arising from the theory of open quantum systems [23–26]; we discuss their decomposability and show how they can be used to detect bound entangled states. In particular, we will be concerned with a class of 16×16 density matrices naturally arising from a 4×4 square lattice through the use of tensor products of Pauli matrices.

In Section 2, we shortly review some basic notions and results concerning positivity, complete positivity and quantum dynamical semigroups; in Section 3, we focus on the above mentioned class of lattice states and study which of them remain positive under partial transposition (PPT); finally, we present some results on the study of their entanglement properties.

2. Positive and completely positive maps

We start with some basic facts about positive maps and entanglement in the case of a finite d-dimensional system S_d. We shall denote by $M_d(\mathbf{C})$ the algebra of $d \times d$ complex matrices and by $\mathcal{S}_d$ the space of states (density matrices) of S_d, that is the convex set of positive $\rho \in M_d(\mathbf{C})$ of unit trace.

Any hermiticity and trace-preserving linear map $\Lambda : \mathcal{S}_d \mapsto \mathcal{S}_d$ can be written as [27]

$$\rho \mapsto \Lambda[\rho] = \sum_{k,i=0}^{d^2-1} \lambda_{ki} \, F_k \, \rho \, F_i^\dagger \ , \tag{1}$$

where the F_k, $k = 1, \ldots, d^2 - 1$, are traceless $d \times d$ matrices, forming together with the normalized identity $F_0 := \mathbf{1}_d/\sqrt{d}$ an orthonormal set in $M_d(\mathbf{C})$: $\mathrm{Tr}\big(F_i^\dagger F_k\big) = \delta_{ki}$, $i, k = 0, 1, \ldots, d^2 - 1$; the coefficients λ_{ki} form a generic hermitian matrix such that $\sum_{k,i=0}^{d^2-1} \lambda_{ki} F_i^\dagger F_k = 1$.

Any linear map Λ that is used to describe a physical state transformation, must preserve the positivity of all states ρ, otherwise the appearance of negative eigenvalues in $\Lambda[\rho]$ would spoil its statistical interpretation, which is based on the use of its eigenvalues as probabilities.

The property of preserving the positivity of the spectrum of all ρ is called positivity; however, it is not sufficient to make Λ fully physically consistent. Indeed, the system S_d may always be thought to be statistically coupled to an ancilla n-level system S_n; one is thus forced to consider the action $\mathrm{id}_n \otimes \Lambda$ over the compound system S_n+S_d, where id_n is the identity action on the ancilla. In order to be fully physically consistent as a state transformation, not only Λ should be positive, but also $\mathrm{id}_n\otimes\Lambda$ for all n; such a property is called complete positivity [3, 7, 8].

Complete positivity is necessary because of the existence of entangled states of the compound system S_n+S_d, namely of states that cannot be written as factorized linear convex combinations

$$\rho^{\mathrm{sep}}_{S_n+S_d}=\sum_{ij}c_{ij}\,\rho^i_{S_n}\otimes\rho^j_{S_d}\ ,\quad c_{ij}>0\ ,\quad \sum_{ij}c_{ij}=1\ . \tag{2}$$

If only separable states as in (2) existed, then positivity of Λ would be enough.

We briefly collect below some results concerning positivity and complete positivity that will be used in the following.

In the space of states $\mathcal{S}_{d\times d}$ of the bipartite system S_d+S_d, let us introduce the symmetric state

$$|\Psi^d_+\rangle=\frac{1}{\sqrt{d}}\sum_{j=1}^{d}|j\rangle\otimes|j\rangle\ , \tag{3}$$

where $|j\rangle$, $j=1,2,\dots,d$ is any fixed orthonormal basis in $\mathbf{C}^d$, and the corresponding projection $P^d_+\equiv|\Psi^d_+\rangle\langle\Psi^d_+|\in\mathcal{S}_{d\times d}$ onto it.

THEOREM 1. *The following results hold:*

(i) Λ *is positive on* $\mathcal{S}_d$ *if and only if*

$$\langle\phi\otimes\psi|\left(\mathrm{id}_d\otimes\Lambda\right)[P^d_+]\,|\phi\otimes\psi\rangle=\frac{1}{d}\langle\psi|\Lambda[\,|\phi^*\rangle\langle\phi^*|\,]|\psi\rangle\geq 0 \tag{4}$$

for all normalized $|\phi\rangle$ *and* $|\psi\rangle\in\mathbf{C}^d$*, with* $|\phi^*\rangle$ *denoting the conjugate of* $|\phi\rangle$ *with respect to the fixed orthonormal basis in* $\mathbf{C}^d$ *[28, 29].*

(ii) Λ *is completely positive on* $\mathcal{S}_d$ *if and only if [3, 30]*

$$\left(\mathrm{id}_d\otimes\Lambda\right)[P^d_+]\geq 0\ . \tag{5}$$

(iii) A linear map Λ *is completely positive on* $\mathcal{S}_d$ *if and only if it can be expressed in the Kraus-Stinespring form [7, 8]*

$$\Lambda[\rho]=\sum_{j=0}^{d^2-1}G_j\,\rho\,G_j^\dagger\ , \tag{6}$$

where the matrices G_j *need not be traceless and must satisfy the condition* $\sum_j G_j^\dagger G_j=1$ *if one asks for trace-preservation.*

REMARK 2. Physically speaking, this Theorem states that, when $P \in \mathcal{S}_{d\times d}$ is a generic one-dimensional projection, positivity of Λ guarantees the positivity of $\big(\mathrm{id}_d \otimes \Lambda\big)[P]$ only when P is separable, *i.e.* $P = |\psi\rangle\langle\psi| \otimes |\phi\rangle\langle\phi|$. Local operations must transform states of the bipartite system $S_d + S_d$ into states keeping the positivity of the associated density matrices; therefore, the existence of entangled states as P^d_+ excludes that Λ, when only positive, may correspond to a physically consistent local state-transformation.

REMARK 3. The positivity or complete positivity of Λ depends on the properties of the $d^2 \times d^2$ matrix of coefficients $[\lambda_{ki}]$ in (1). Indeed, the Kraus-Stinespring form of completely positive maps corresponds to $[\lambda_{ki}] = \mathrm{diag}(1,1,\ldots,1)$. Vice versa, if the matrix $[\lambda_{ki}]$ is positive, diagonalizing $\lambda_{ki} = \sum_{j=0}^{d^2-1} \ell_j U_{kj} U^*_{ij}$ and setting $G_j = \sum_{k=0}^{d^2-1} \sqrt{\ell_k} U_{kj} F_k$, one recovers (6).

REMARK 4. The symmetric state $|\Psi^d_+\rangle$ and the corresponding density matrix P^d_+ satisfy the following properties:

(i) for all matrices A, B acting on $\mathbf{C}^d$ one has

$$A \otimes B |\Psi^d_+\rangle = \mathbf{1}_d \otimes BA^T |\Psi^d_+\rangle = AB^T \otimes \mathbf{1}_d |\Psi^d_+\rangle \ , \tag{7}$$

where A^T, B^T denote the transposed of A, B;

(ii) under partial transposition, P^d_+ gives rise to the flip operator

$$V = \sum_{a,b=1}^{d} |a\rangle\langle b| \otimes |b\rangle\langle a| \ = \ d\,\big(T_d \otimes \mathrm{id}_d\big)[P^d_+] \ , \tag{8}$$

which is such that

$$V|\psi \otimes \phi\rangle = |\phi \otimes \psi\rangle \ , \quad V\big(A \otimes B\big)V = B \otimes A \ . \tag{9}$$

Unlike for completely positive maps, there is no general prescription on $[\lambda_{ki}]$ ensuring that Λ preserve the positivity of ρ. For instance, if $[\lambda_{ki}]$ is not positive, then, by separating positive and negative eigenvalues, one sees that every Λ can be written as the difference of two completely positive maps:

$$\Lambda[\rho] = \sum_{\ell_j \geq 0} \ell_j \, G_j \, \rho \, G_j^\dagger \ - \ \sum_{\ell_j < 0} |\ell_j| \, G_j \, \rho \, G_j^\dagger \ , \tag{10}$$

where the matrices $G_j := \sum_{k=0}^{d^2-1} U_{kj} F_k$, like the F_k's, are orthogonal. However, no general rule is known that may allow us to recognize the positivity of Λ by looking at the eigenvalues ℓ_k and at the matrices G_k. One has to content oneself with sufficient conditions as the one which follows, that assumes the existence of just one negative eigenvalue.

PROPOSITION 5. *Suppose that $\ell_k \geq 0$ for all $k \neq p$, while $\ell_p < 0$, whence (10) reads*

$$\Lambda[\rho] = \sum_{\ell_k \geq 0} \ell_k \, G_k \, \rho \, G_k^\dagger \; - \; |\ell_p| \, G_p \, \rho \, G_p^\dagger \; . \tag{11}$$

If $\|G_p\|^2 = M < 1$ and $\ell_k \geq \frac{M}{1-M}|\ell_p|$, $k \neq p$, then Λ is positive.

The condition (4) is satisfied since

$$\langle\psi|\left(\sum_{k=0}^{d^2-1} \ell_k \, G_k|\phi^*\rangle\langle\phi^*|G_k^\dagger\right)|\psi\rangle \geq |\ell_p|\left(\frac{M}{1-M}\sum_{k\neq p}|\langle\psi|G_k|\phi^*\rangle|^2 - |\langle\psi|G_p|\phi^*\rangle|^2\right)$$
$$= |\ell_p| \, \frac{M}{1-M}\left(1 - \frac{1}{M} \, |\langle\psi|G_p|\phi^*\rangle|^2\right) \geq 0 \; ,$$

where the last equality holds because the G_k's form a basis in $M_d(\mathbf{C})$, whence

$$\sum_{k=0}^{d^2-1} |\langle\psi|G_k|\phi^*\rangle|^2 = \mathrm{Tr}\left[\left(\sum_{k=0}^{d^2-1} G_k \, \mathrm{Tr}\left[G_k^\dagger|\psi\rangle\langle\phi^*|\right]\right)|\phi^*\rangle\langle\psi|\right] = \langle\phi^*|\phi^*\rangle\langle\psi|\psi\rangle = 1 \; .$$

■

EXAMPLE 6. Let $d = 2$ and σ_α, $\alpha = 0,1,2,3$, be the Pauli matrices and the 2×2 identity matrix σ_0. Let $S_\alpha : \mathcal{S}_2 \mapsto \mathcal{S}_2$ denote the positive map $\rho \mapsto \sigma_\alpha\rho\sigma_\alpha$, and consider the maps

$$\rho \mapsto \mathrm{Tr}_2[\rho] := \frac{1}{2}\sum_{\alpha=0}^{3} S_\alpha[\rho] \; , \quad \rho \mapsto T_2[\rho] := \frac{1}{2}\sum_{\alpha=0}^{3} \varepsilon_\alpha S_\alpha[\rho] \; , \tag{12}$$

where $\varepsilon_\alpha = 1$ when $\alpha \neq 2$, whereas $\varepsilon_2 = -1$. By developing $\rho = \sum_{\alpha=0}^{3} \rho_\alpha \, \sigma_\alpha$, it is straightforward to check that the first map amounts to taking the trace and multiplying it by the identity σ_0; it is completely positive: indeed, by taking $F_\alpha = \sigma_\alpha/\sqrt{2}$ in (1), one finds $[\lambda_{\alpha\beta}] = \mathrm{diag}(1,1,1,1)$. On the other hand, T_2 corresponds to transposition and is only positive, for $[\lambda_{\alpha\beta}] = \mathrm{diag}(1,1,-1,1)$ and $\|F_\alpha\|^2 = 1/2$.

As observed in Remark 2, a positive map Λ that is not completely positive is unphysical because its "extension" by the identity map, *i.e.* $\mathrm{id}_d \otimes \Lambda$, moves entangled states as P_d^+ out of the space of states. However, exactly because of this it may be used to detect entanglement [15].

THEOREM 7. *A state $\rho \in \mathcal{S}_{d\times d}$ is entangled iff*

$$D_\Lambda[\rho] := \mathrm{Tr}\Big[\big(\mathrm{id}_d \otimes \Lambda\big)[P_+^d] \, \rho\Big] \; < \; 0 \tag{13}$$

for some positive map Λ on $\mathcal{S}_d$.

Indeed, notice that according to (4), $D_\Lambda[\rho^{\rm sep}] \geq 0$ for any separable state $\rho^{\rm sep}$.

REMARK 8. For any map Λ, its dual Λ^* is defined by $\mathrm{Tr}(\Lambda[X]\rho) = \mathrm{Tr}(X\Lambda^*[\rho])$, for any ρ, X. Then, the above result can be formulated by saying that $\rho \in \mathcal{S}_{d\times d}$ is entangled if and only if there exists a positive map Λ on $\mathcal{S}_d$ such that $\big(\mathrm{id}_d \otimes \Lambda^*\big)[\rho]$ is non-positive.

As mentioned in the Introduction, an important example of positive, but not completely positive map is the transposition T_d (w.r.t. a fixed basis in $\mathbf{C}^d$); through it, one can construct a subset of positive maps, the cone of decomposable maps.

DEFINITION 9. A positive map Λ on $\mathcal{S}_d$ is decomposable if

$$\Lambda = \Gamma_1 + \Gamma_2 \circ T_d \ , \tag{14}$$

with $\Gamma_{1,2}$ completely positive on $\mathcal{S}_d$.

REMARK 10. If $\big(\mathrm{id}_d \otimes T_d\big)[\rho]$ is positive, then $\big(\mathrm{id}_d \otimes \Lambda\big)[\rho]$ is also positive for any decomposable Λ. Thus, if $\rho \in \mathcal{S}_{d\times d}$ is entangled, but with positive partial transpose (PPT), a decomposable Λ can not detect its entanglement since $D_\Lambda(\rho) \geq 0$.

As a consequence, one has the following result.

PROPOSITION 11. *If Λ is positive on $\mathcal{S}_d$, $\rho \in \mathcal{S}_{d\times d}$ is PPT and $D_\Lambda(\rho) < 0$, then Λ is not decomposable and ρ is entangled.*

For $d = 2$, all positive maps result decomposable [31], whence the transposition detects all the entangled states. In other words, $\big(\mathrm{id}_2 \otimes T_2\big)[\rho]$ is non-positive if and only if ρ is entangled. On the contrary, when $d \geq 3$, there are PPT states which are entangled [4,6,16,18]; their entanglement can not be distilled by means of local operations and classical communication [17], and it is called bound entanglement. It is thus important not only to identify entanglement when present, but also to qualify whether it is bound or not.

2.1. POSITIVE AND COMPLETELY POSITIVE SEMIGROUPS

The unitary time evolution for a system S_d generated by a Hamiltonian operator results automatically completely positive due to the criterion (iii) in Theorem 1. On the other hand, when the system is in contact with an external environment, its subdynamics is in general not unitary, incorporating noise and dissipative effects. In many physically interesting cases, such dynamics can be approximated by families of linear maps γ_t, $t \geq 0$ on $\mathcal{S}_d$ that are not invertible and obey a semigroup composition law, $\gamma_t \circ \gamma_s = \gamma_{t+s}$ [23–26].

Under the mild assumptions of continuity in time, $\lim_{t\to 0}\gamma_t=\mathrm{id}_d$, and preservation of hermiticity and trace, the semigroup has the form $\gamma_t=\exp(tL)$ with generator [27]

$$L[\rho]=-i\Big[H\,,\,\rho_t\Big]\,+\sum_{i,j=1}^{d^2-1}C_{ij}\Big(F_i\rho_tF_j^\dagger\,-\,\frac{1}{2}\Big\{F_j^\dagger F_i\,,\,\rho_t\Big\}\Big)\ , \tag{15}$$

where the F_j are as in (1), while the matrix of coefficients $[C_{ij}]$, called the Kossakowski matrix, is hermitian.

Regarding the generated maps $\exp(tL)$, there are no general results on the form of the Kossakowski matrix $[C_{ij}]$ such that they are positive on $\mathcal{S}_d$, whereas they are completely positive if and only if $[C_{ij}]\geq 0$.

In the following we shall be interested in semigroups $\Gamma_t=\gamma_t^1\otimes\gamma_t^2$ on $\mathcal{S}_{d\times d}$ that are tensor products of semigroups $\gamma_t^{1,2}=\exp(tL_{1,2})$ on $\mathcal{S}_d$. Like in Proposition 5, one can give simple conditions on the Kossakowski matrices $C_{1,2}$ in the generators $L_{1,2}$ that are sufficient for the positivity of Γ_t [22].

PROPOSITION 12. *Suppose that the non-Hamiltonian terms in the generators of $\gamma_t^{1,2}$ are as follows,*

$$L_i[\rho]=\sum_{\ell=1}^{d^2-1}c_i^\ell\Big(G^i_\ell\,\rho\,G^i_\ell\,-\,\frac{1}{2}\Big\{(G^i_\ell)^2\,,\,\rho\Big\}\Big)\ ,\quad c_i^\ell\in\mathbf{R}\ , \tag{16}$$

for $i=1,2$, where $G^i_\ell\in M_d(\mathbf{C})$, together with $G^i_0=\mathbf{1}_2/\sqrt{d}$, constitute two orthonormal sets of hermitian matrices. Suppose that $c_1^\ell>0$ for all $\ell=1,2,\ldots,d^2-1$, and that $c_2^k=-|c_2^k|<0$, for one index k, while $c_2^\ell>0$ when $\ell\neq k$; then, the map $\Gamma_t=\gamma_t^1\otimes\gamma_t^2$ is positive if $c_1^\ell\geq|c_2^k|$, $\ell=1,2,\ldots,d^2-1$ and $c_2^\ell\geq|c_2^k|$, $\ell=1,2,\ldots,d^2-1$, $\ell\neq k$.

EXAMPLE 13. Let $d=2$, set $H=0$, $F_i=\sigma_i/\sqrt{2}$ and choose as Kossakowski matrices

$$C_1=\begin{pmatrix}1&0&0\\0&1&0\\0&0&1\end{pmatrix}\ ,\qquad C_2=\begin{pmatrix}1&0&0\\0&-1&0\\0&0&1\end{pmatrix}\ . \tag{17}$$

Using the notation of Example 6, the generators are given by

$$\partial_t\gamma_t^1[\rho]=L_1[\rho]:=\frac{1}{2}\Big(\sum_{i=1}^3S_i[\rho]\,-\,3\,\rho\Big)\ ,\quad\partial_t\gamma_t^2[\rho]=L_2[\rho]:=\frac{1}{2}\Big(\sum_{i=1}^3\varepsilon_i\,S_i[\rho]\,-\,\rho\Big)\ , \tag{18}$$

yielding

$$\gamma_t^1=\mathrm{e}^{-2t}\,\mathrm{id}_2+\frac{1-\mathrm{e}^{-2t}}{2}\,\mathrm{Tr}_2\ ,\quad\gamma_t^2=\frac{1+\mathrm{e}^{-2t}}{2}\,\mathrm{id}_2+\frac{1-\mathrm{e}^{-2t}}{2}T_2\ . \tag{19}$$

As $T_4 = T_2 \otimes T_2$ and $\mathrm{Tr}_2 \circ T_2 = \mathrm{Tr}_2$, $\Gamma_t = \gamma_t^1 \otimes \gamma_t^2$ can be written as

$$\begin{aligned}\Gamma_t &= \mathrm{e}^{-2t}\frac{1+\mathrm{e}^{-2t}}{2}\,\mathrm{id}_4 + \frac{1-\mathrm{e}^{-4t}}{4}\,\mathrm{Tr}_2 \otimes \mathrm{id}_2 \\ &+ \frac{1-\mathrm{e}^{-2t}}{2}\Big(\mathrm{e}^{-2t}T_2 \otimes \mathrm{id}_2 + \frac{1-\mathrm{e}^{-2t}}{2}\,\mathrm{Tr}_2 \otimes \mathrm{id}_2\Big) \circ T_4 \ . \qquad (20)\end{aligned}$$

The semigroup Γ_t is written as $\Gamma_t = \Gamma_t^1 + \Gamma_t^2 \circ T_4$, with Γ_t^1 completely positive for all $t \geq 0$, while Γ_t^2 is such only for $t = 0$ and $t \geq t^* := (\log 3)/2$. It follows that Γ_t is surely decomposable for $t = 0$ ($\Gamma_0 = \mathrm{id}_{16}$) and for $t \geq t^*$; on the other hand, as shown below the positive maps Γ_t result indecomposable for $0 < t < t^*$.

3. Lattice states

Let L_{16} be the square lattice $\big\{(\alpha,\beta)\,:\,\alpha\,,\,\beta = 0,1,2,3\big\}$ with 16 elements, that we can split into columns $\mathcal{C}_\alpha = \big\{(\alpha,\beta)\,:\,\beta = 0,1,2,3\big\}$ and rows $\mathcal{R}_\beta = \big\{(\alpha,\beta)\,:\,\alpha = 0,1,2,3\big\}$. To each site (α,β) we will associate the following 4×4 tensor products of Pauli matrices $\sigma_{\alpha\beta} := \sigma_\alpha \otimes \sigma_\beta$. By acting with $\mathbf{1}_4 \otimes \sigma_{\alpha\beta}$ on the symmetric state $|\Psi_+^4\rangle$ one constructs an orthonormal basis of maximally entangled vectors in $\mathbf{C}^{16}$,

$$|\Psi_{\alpha\beta}\rangle := \big(\mathbf{1}_4 \otimes \sigma_{\alpha\beta}\big)\,|\Psi_+^4\rangle, \qquad (21)$$

with the corresponding orthogonal projections,

$$P_{\alpha\beta} := |\Psi_{\alpha\beta}\rangle\langle\Psi_{\alpha\beta}| = \big(\mathbf{1}_4 \otimes \sigma_{\alpha\beta}\big)\,P_+^4\,\big(\mathbf{1}_4 \otimes \sigma_{\alpha\beta}\big)\ , \qquad P_{\alpha\beta}\,P_{\gamma\varepsilon} = \delta_{\alpha\gamma}\,\delta_{\beta\varepsilon}\,P_{\alpha\beta}\ . \quad (22)$$

We will focus on states in $\mathcal{S}_4 \times \mathcal{S}_4$ that are diagonal with respect to the just introduced basis $\big\{|\Psi_{\alpha\beta}\rangle\big\}$,

$$\rho_\pi := \sum_{(\alpha,\beta)\in L_{16}} \pi_{\alpha\beta}\,P_{\alpha\beta}\ , \quad \pi_{\alpha\beta} \geq 0\ , \quad \sum_{(\alpha\beta)\in L_{16}} \pi_{\alpha\beta} = 1\ . \qquad (23)$$

Actually, for sake of simplicity we shall further limit the discussion to those states ρ_I for which all the non-vanishing weights $\pi_{\alpha\beta}$ are equal. They are completely characterized by the subset $I = \{(\alpha,\beta) : \pi_{\alpha\beta} \neq 0\} \subseteq L_{16}$, and take the form

$$\rho_I = \frac{1}{N_I}\sum_{(\alpha,\beta)\in I} P_{\alpha\beta}\ , \qquad (24)$$

where N_I is the number of elements of I.

We shall now characterize all states ρ_I that are PPT and study their entanglement properties.

3.1. PPT States

We first act with the transposition on the first subsystem in (23):

$$\rho_\pi^{T_1} := \big(T_4\otimes \mathrm{id}_4\big)[\rho_\pi] = \frac{1}{4} \sum_{(\alpha,\beta)\in L_{16}} \pi_{\alpha\beta}\, V_{\alpha\beta}\ , \qquad V_{\alpha\beta} := \big(\mathbf{1}_4\otimes\sigma_{\alpha\beta}\big)\, V\, \big(\mathbf{1}_4\otimes\sigma_{\alpha\beta}\big)\ . \tag{25}$$

LEMMA 14. *The matrices* $V_{\alpha\beta}$ *are self-adjoint and their spectral decomposition is as follows:* $V_{\alpha\beta} = \sum_{(\gamma,\delta)\in L_{16}} \xi_{\alpha\gamma}\,\xi_{\beta\delta}\,P_{\gamma\delta}$ *where* $\xi_{\alpha\gamma} = 1\,-\,2\,\delta_{|\alpha-\gamma|,2}$.

From (9) and $V|\Psi_+^4\rangle = |\Psi_+^4\rangle$, it follows that

$$\begin{aligned} V_{\alpha\beta}|\Psi_{\gamma\delta}\rangle &= \varepsilon_\alpha\varepsilon_\gamma\varepsilon_\beta\varepsilon_\delta\Big\{\mathbf{1}_4\otimes\Big[\big(\sigma_\alpha\sigma_\gamma\sigma_\alpha\big)\otimes\big(\sigma_\beta\sigma_\delta\sigma_\beta\big)\Big]\Big\}|\Psi_+^4\rangle \\ &= \big(\varepsilon_\alpha\varepsilon_\gamma\eta_{\alpha\gamma}\big)\ \big(\varepsilon_\beta\varepsilon_\delta\eta_{\beta\delta}\big)\ |\Psi_{\gamma\delta}\rangle\ , \end{aligned}$$

where $\eta_{\alpha\gamma}$ takes the values ±1 according to the table

$\gamma\backslash\alpha$	0	1	2	3
0	1	1	1	1
1	1	1	-1	-1
2	1	-1	1	-1
3	1	-1	-1	1

and ε_α has been defined in Example 6. Setting $\xi_{\alpha\gamma} = \varepsilon_\alpha\varepsilon_\gamma\,\eta_{\alpha\gamma}$, the result follows by direct inspection. ■

PROPOSITION 15. *A state* ρ_I *is PPT if and only if for any lattice point* $(\alpha,\beta)\in L_{16}$ *the corresponding column* $\mathcal{C}_\alpha$ *and row* $\mathcal{R}_\beta$ *do not contain more than* $N_I/2$ *elements of* I*, the point* (α,β) *itself excluded.*

It proves convenient to consider the convex set of states ρ_π introduced in (23). Recalling the explicit form of the coefficients $\xi_{\alpha\gamma}$ above, we introduce the bijection $(\alpha,\beta)\mapsto(\widetilde{\alpha},\widetilde{\beta})$, where $\widetilde{\mu} := (\mu+2)\ \mathrm{mod}\,(4)$, $\mu = 0,\ldots,3$. Then, from (25), using the previous Lemma, we obtain the spectral decomposition

$$\rho_\pi^{T_1} = \sum_{(\gamma,\delta)\in L_{16}} \frac{1}{4}\Big[1 - 2\Big(Q^{(1)}_{\widetilde{\gamma}\widetilde{\delta}} + Q^{(2)}_{\widetilde{\gamma}\widetilde{\delta}}\Big)\Big]\, P_{\gamma\delta} \tag{26}$$

$$Q^{(1)}_{\widetilde{\gamma}\widetilde{\delta}} := \sum_{\beta\neq\widetilde{\delta}} \pi_{\widetilde{\gamma}\beta}\ , \qquad Q^{(2)}_{\widetilde{\gamma}\widetilde{\delta}} := \sum_{\alpha\neq\widetilde{\gamma}} \pi_{\alpha\widetilde{\delta}}\ , \tag{27}$$

whence ρ_π is PPT if and only if $Q^{(1)}_{\widetilde{\gamma}\widetilde{\delta}} + Q^{(2)}_{\widetilde{\gamma}\widetilde{\delta}} \leq \dfrac{1}{2}$ for all $(\gamma, \delta) \in L_{16}$. Because of the bijection $(\mu, \nu) \mapsto (\widetilde{\mu}, \widetilde{\nu})$, one has, equivalently, that ρ_π is PPT if and only if $Q^{(1)}_{\gamma\delta} + Q^{(2)}_{\gamma\delta} \leq \dfrac{1}{2}$ for all $(\gamma, \delta) \in L_{16}$. Setting $\pi_{\alpha\beta} = 1/N_I$ on $I \subseteq L_{16}$ and $\pi_{\alpha\beta} = 0$ otherwise, the result follows, for the quantities $Q^{(1,2)}$ just count how many contributions come from any given row and column of the lattice, the intersection point (α, β) excluded. ■

REMARK 16. In order to concretely construct the states ρ_I that are PPT, one can proceed by first selecting those subsets $I \subseteq L_{16}$ containing a decreasing number of elements in the ordered columns $\mathcal{C}_0$, $\mathcal{C}_1$, $\mathcal{C}_2$, $\mathcal{C}_3$, with those in $\mathcal{C}_0$ piled one over the other, without holes, and further satisfying the hypothesis of Proposition 15.

EXAMPLE 17. Representing the elements of I by crosses inserted at the corresponding sites of the lattice L_{16}, one can associate to each state ρ_I a graph. Using this correspondence, we give below some examples PPT states.

$N_I = 4$:

3				×
2			×	
1		×		
0	×			
	0	1	2	3

3				
2				
1	×	×		
0	×	×		
	0	1	2	3

$N_I = 6$:

3				
2				
1	×	×	×	
0	×	×	×	
	0	1	2	3

3				
2	×			×
1	×	×		
0	×		×	
	0	1	2	3

$N_I = 8$:

3				
2	×	×	×	
1	×	×		
0	×	×	×	
	0	1	2	3

3				
2	×	×		×
1	×	×		
0	×	×	×	
	0	1	2	3

$N_I = 10$:

3				
2	×	×	×	×
1	×	×		×
0	×	×	×	
	0	1	2	3

3				
2	×	×	×	×
1	×	×	×	
0	×	×	×	
	0	1	2	3

In addition, it is instructive to give some examples of states whose partial transpose is not positive; in each of them the condition of Proposition 15 is not satisfied, as easily seen by considering the highlighted columns and rows.

$N_I = 4$:



3				
2		×		
1	×			
0	×	×		
	0	1	2	3

$N_I = 5$:



3				
2				
1	×	×		
0	×	×	×	
	0	1	2	3

$N_I = 6$:



3				
2			×	
1	×	×		
0	×	×	×	
	0	1	2	3

$N_I = 7$:



3				
2				
1	×	×	×	
0	×	×	×	×
	0	1	2	3

All lattice states ρ_I that result PPT can be obtained by those defined in Remark 16 by applying suitable local unitary transformations. Indeed, one has:

PROPOSITION 18. *Let U and W be 4×4 unitary matrices, transforming $\sigma_{\alpha\beta} \mapsto W\sigma_{\alpha\beta}U^\dagger$ into another element $\sigma_{\gamma\delta}$ up to a phase. Then, any PPT state ρ_I is mapped into another PPT state ρ_I of the same rank,*

$$\rho_{\hat{I}} := \left(U^* \otimes W\right) \rho_I \left(U^T \otimes W^\dagger\right) \ , \tag{28}$$

where $\hat{I} \subseteq L_{16}$ is the image of I under the one-to-one correspondence $(\alpha,\beta) \mapsto (\gamma,\delta)$.

The result follows at once from the definition of $P_{\alpha\beta}$, the property (7) of completely symmetric states and the hypothesis; in fact

$$\begin{aligned} \left(U^* \otimes W\right) P_{\alpha\beta} \left(U^T \otimes W^\dagger\right) &= \Big[\mathbf{1}_4 \otimes \left(W\sigma_{\alpha\beta}U^\dagger\right)\Big] |\Psi^4_+\rangle\langle\Psi^4_+| \Big[\mathbf{1}_4 \otimes \left(U\sigma_{\alpha\beta}W^\dagger\right)\Big] \\ &= P_{\gamma\delta} \ , \end{aligned}$$

and the local unitary operations preserve the PPT property. ■

By properly choosing U and W, one can permute columns and rows. Further, by taking U and W both equal to the flip operator V in (8), one can go from any ρ_I to the one with column and row contributions exchanged. More in general, one can subdivide the PPT states ρ_I into equivalence classes with respect to the relation given by (28).

From the previous considerations, by direct inspection one can conclude that for $N_I = 1, 2, 3, 5, 7$, there are no states ρ_I that are PPT, while they are all PPT for $N_I \geq 12$.

3.2. Bound Entangled States

Having fully characterized the states ρ_I that are PPT, the next task is to determine which of them are also entangled. Unfortunately, we are not able to obtain a complete classification. Nevertheless, by employing the semigroup of positive maps Γ_t constructed in the previous section we can discuss the entanglement properties of some non-trivial classes of PPT states ρ_I.

We first observe that:

(i) from general arguments rank-4 states cannot be entangled [32];

(ii) all PPT lattice states ρ_I, with $N_I = 15$, are separable isotropic states; indeed, thanks to Proposition 18, all of them are in the equivalence class of $\rho_{15} = (1 - P_{00})/15$, with fidelity $\mathcal{F} = \mathrm{Tr}[\rho_{15}\, P_{00}] = 0$ (*cf.* [30]).

Further, we recall the following result discussed in [22].

EXAMPLE 19. Recall the definition of the pairing D_Λ in (13) and take for Λ the map Γ_t as in (20); consider the following state $\rho_I \in \mathcal{S}_4$, that according to the rule in Proposition 15 is PPT

$$\rho_I = \frac{1}{6}\Big(P_{02} + P_{11} + P_{23} + P_{31} + P_{32} + P_{33}\Big)\ . \tag{29}$$

It turns out that $D_{\Gamma_t}(\rho_I) < 0$ for $0 < t < (\ln 3)/2$. As a consequence, ρ is bound entangled and, in this range of times, the positive maps Γ_t turn out to be indecomposable.

Actually, the positive semigroup Γ_t is able to detect the entanglement of other PPT lattice states ρ_I. Instead of using the test (13), we will follow a different, equivalent method, based on Remark 8: if for a PPT state ρ_I, any of the eigenvalues of $\mathrm{id}_4 \otimes \Gamma_t[\rho_I]$ is negative, then we can conclude that ρ_I is bound entangled (notice that the dual of Γ_t coincides with Γ_t itself).

It is straightforward to check that

$$\langle\Psi_{\gamma\delta}|\big(\,\mathrm{id}_4 \otimes \Gamma_t\big)[P_{\alpha\beta}]|\Psi_{\mu\nu}\rangle = \delta_{\gamma\mu}\,\delta_{\delta\nu}\ p^{\mu\nu}_{\alpha\beta}(t)\ ,$$

with

$$p^{\mu\nu}_{\alpha\beta}(t) := \frac{1}{16}\Big(4\mathrm{e}^{-2t}\,\delta_{\alpha\mu}\,+\,1\,-\,\mathrm{e}^{-2t}\Big)\,\times\,\Big[2\Big(1+\mathrm{e}^{-2t}\Big)\delta_{\beta\nu}\,+\,\Big(1-\mathrm{e}^{-2t}\Big)\xi_{\beta\nu}\Big]\ , \tag{30}$$

where $\xi_{\beta\delta}$ has been introduced in Lemma 14. Therefore, we can write the following spectral decomposition

$$\big(\mathrm{id}_4 \otimes \Gamma_t\big)[\rho_I] = \sum_{(\mu,\nu)\in L_{16}} R_{\mu\nu}(t)\ P_{\mu\nu}\ , \quad R_{\mu\nu}(t) := \frac{1}{N_I} \sum_{(\alpha,\beta)\in I} p^{\mu\nu}_{\alpha\beta}(t)\ . \tag{31}$$

If one of the eigenvalues $R_{\mu\nu}(t) < 0$, the PPT state ρ_I is entangled and its entanglement of bound type and thus undistillable.

PROPOSITION 20. *A PPT lattice state ρ_I is entangled if there exist a column $\mathcal{C}_\gamma$ and a row $\mathcal{R}_\delta$ of L_{16} whose intersection with the subset I contains only one element, (γ, δ'), $\delta' \neq \delta$, or (γ', δ), $\gamma' \neq \gamma$.*

By expanding the eigenvalues of $\big(\mathrm{id}_4 \otimes \Gamma_t\big)[\rho_I]$ for small t, one gets

$$p^{\mu\nu}_{\alpha\beta}(t) \simeq (1-3t)\delta_{\alpha\mu}\delta_{\beta\nu}\ +\ \frac{t}{2}\Big(\delta_{\alpha\mu}\,\xi_{\beta\nu}\ +\ \delta_{\beta\nu}\Big)\ . \tag{32}$$

Let us suppose that the only element of I in the column $\mathcal{C}_\gamma$ and the row $\mathcal{R}_\delta$ be (γ, δ'), $\delta' \neq \delta$. According to Proposition 18, we can always permute rows and columns so that $\gamma = \delta = 3$, and $\delta' = 1$, while remaining in the same equivalence class. According to the assumptions and to the definition of ξ_{ν_β} in Lemma 14, it turns out that, for small positive t, the eigenvalue R_{33} is negative:

$$R_{33}(t) = \frac{1}{N_I} \sum_{(\alpha,\beta)\in I} p^{33}_{\alpha\beta}(t)\ =\ -\frac{t}{2N_I}\ . \tag{33}$$

as it can be easily checked by inserting (32) into (31). ■

Notice that the previous Proposition gives only a sufficient condition for a state ρ_I to be entangled and further that its hypothesis can be satisfied only when $N_I \leq 10$. Nevertheless, using it we are able to conclude that among the PPT cases presented in Example 17, with $6 \leq N_I \leq 10$, those on the right result entangled. On the other hand, Proposition 20 is inconclusive for what concerns the states on the left; indeed, in those case, $\big(\mathrm{id}_4 \otimes \Lambda\big)[\rho_I]$ with $\Lambda = \Gamma_t$ has no negative eigenvalues for small t, but these might occur for a different Λ.

Actually, we have been unable to find a bound entangled lattice state ρ_I that is not detected by the positive map Γ_t; this fact may suggest that Γ_t is sufficient to classify all entangled PPT state, within the studied class.

For instance, the rank-6 PPT state on the left in Example 17 results separable. In order to prove this, first use Proposition 18 to transform it into the following element of its equivalence class:

$$\rho_6 = \frac{1}{6}\left(P_{00} + P_{01} + P_{02} + P_{30} + P_{31} + P_{32}\right)\ .$$

It is then lengthy, but not difficult to decompose

$$\begin{aligned}\rho_6 &= \frac{1}{12}\Big[P^{(1)}_{0+}+P^{(1)}_{0-}+P^{(1)}_{1+}+P^{(1)}_{1-}+P^{(2)}_{0+}+P^{(2)}_{0-}+P^{(2)}_{1+}+P^{(2)}_{1-}\Big]\\ &+ \frac{1}{12}\Big[Q_3+Q_4+Q_5+Q_6\Big]\ ,\end{aligned}$$

where the P's and Q's are projections onto the separable states:

$$\begin{aligned}|\Psi^{(1)}_{0\pm}\rangle &= \frac{|00\rangle \pm |01\rangle}{\sqrt{2}}\otimes\frac{|00\rangle \pm |01\rangle}{\sqrt{2}} \qquad |\Psi^{(2)}_{0\pm}\rangle = \frac{|00\rangle \pm i|01\rangle}{\sqrt{2}}\otimes\frac{|00\rangle \mp i|01\rangle}{\sqrt{2}}\ ,\\ |\Psi^{(1)}_{1\pm}\rangle &= \frac{|10\rangle \pm |11\rangle}{\sqrt{2}}\otimes\frac{|10\rangle \pm |11\rangle}{\sqrt{2}} \qquad |\Psi^{(2)}_{1\pm}\rangle = \frac{|10\rangle \pm i|11\rangle}{\sqrt{2}}\otimes\frac{|10\rangle \mp i|11\rangle}{\sqrt{2}}\ ,\end{aligned}$$

$$\begin{aligned}|\Phi_3\rangle &= |00\rangle\otimes|01\rangle \qquad |\Phi_4\rangle = |01\rangle\otimes|00\rangle\ ,\\ |\Phi_5\rangle &= |10\rangle\otimes|11\rangle \qquad |\Phi_6\rangle = |11\rangle\otimes|10\rangle\ ,\end{aligned}$$

respectively, with $|0\rangle$ and $|1\rangle$ eigenstates of σ_3.

Another example is given by the already mentioned rank-15 states, but the same conclusion holds also for rank-14 states. Indeed, all these latter states are in a sigle equivalence class, whose representative element is given by

3	○	○		
2	○	○	×	×
1	◇	◇	×	×
0	◇	◇	×	×
	0	1	2	3

The diamonds, circles and crosses identify lattice states of rank four and six that we have already shown to be separable and whose linear convex combination gives the chosen rank-14 lattice state.

We conclude by mentioning that most of the results presented here can be generalized to the case of higher dimensional bipartite systems. In particular, one can construct $2^N \times 2^N$-dimensional states whose bound entanglement is detected by dynamical semigroups of positive, non-decomposable maps that generalize the map Γ_t discussed in Example 13 [33].

Bibliography

1. M.A. Nielsen and I.L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, Cambridge, 2000.
2. *Quantum Information: An Introduction to Basic Theoretical Concepts and Experiments*, G. Alber et al. eds, Springer Tracts in Modern Physics, **173**, Springer, Berlin, 2001.

3. M. D. Choi, Canad. J. Math. **24**, 520 (1972).
4. E. Störmer, Proc. Amer. Math. Soc. **86**, 402 (1982).
5. K. Życzkowski and I. Bengtsson, Open Sys. & Information Dyn. **11**, 3 (2004).
6. M.D. Choi, Linear Alg. Appl. **10**, 285 (1975).
7. M. Takesaki, *Theory of operator algebras*, Vol. 1, Springer, New York 1979.
8. K. Kraus, *States, Effects and Operations: Fundamental Notions of Quantum Theory*, Lec. Notes Phys. **190**, Springer, Berlin 1983
9. F. Benatti, R. Floreanini, Banach Centre Publications **43**, 71 (1998).
10. F. Benatti, R. Floreanini, R. Romano, J. Phys. A **35**, L551 (2002).
11. F. Benatti, R. Floreanini, R. Romano, J. Phys. A **35**, 4955 (2002).
12. F. Benatti, R. Floreanini, M. Piani, Phys. Rev. A **67**, 042110 (2003).
13. F. Benatti, R. Floreanini, M. Piani and R. Romano, Complete positivity and dissipative factorized dynamics: some comments, *Proceedings of the IV Workshop on Time Asymmetric Quantum Mechanics*, Lisboa, Portugal, 2003, `quant-ph/0310151`.
14. A. Peres, Phys. Rev. Lett. **77**, 1413 (1996).
15. M. Horodecki, P. Horodecki and R. Horodecki, Phys. Lett. A **223**, 1 (1996).
16. P. Horodecki, Phys. Lett. A **232**, 333 (1997).
17. M. Horodecki, P. Horodecki and R. Horodecki, Phys. Rev. Lett. **80**, 5239 (1998).
18. M. Horodecki, P. Horodecki and R. Horodecki, *Mixed-state entanglement and quantum communication*, in: [2].
19. M.D. Choi, Linear Alg. Appl. **12**, 95 (1975).
20. K.-C. Ha, S.-H. Kye and Y.-S. Park, Phys. Lett. A **313**, 163 (2003).
21. A. Kossakowski, Open Sys. and Inf. Dyn. **10**, 1 (2003).
22. F. Benatti, R. Floreanini and M. Piani, Phys. Lett. A **326**, 187 (2004).
23. H. Spohn, Rev. Mod. Phys. **52**, 569 (1980).
24. R. Alicki, K. Lendi, *Quantum Dynamical Semigroups and Applications*, Lec. Notes Phys. **286**, Springer, Berlin 1987.
25. H.-P. Breuer, F. Petruccione, *Theory of Open Quantum Systems*, Oxford University Press, Oxford 2002.
26. *Dissipative Quantum Dynamics*, F. Benatti and R. Floreanini eds., Lec. Notes Phys. **612**, Springer, Berlin 2003.
27. V. Gorini, A. Kossakowski and E. C. G. Sudarshan, J. Math. Phys. **17**, 821 (1976).
28. A. Jamiolkowski, Rep. Math. Phys. **3**, 275 (1972).
29. A. Kossakowski, Bull. Acad. Pol. Sc. **12**, 1021 (1972).
30. M. Horodecki and P. Horodecki, Phys Rev A **59**, 4206 (1999).
31. S.L. Woronowicz, Rep. Math. Phys. **10**, 165 (1976).
32. P. Horodecki, M. Lewenstein, G. Vidal and I. Cirac, Phys Rev A **62**, 032310 (2000).
33. M. Piani, *A class of* $2^N \times 2^N$ *bound entangled states detected by non-decomposable maps*, in preparation